PROJECT TITLE: Winter Wheat Variety Evaluations

YEAR/PROJECT: 1989/756

PROJECT PERSONNEL: Leader - Vern R. Stewart, Todd K. Keener - Research

Specialist, NWARC, Kalispell, MT

SUMMARY:

Severe winter temperatures destroyed the Intrastate and Western Regional Soft White winter wheat nurseries this season. Because of the winter injury, stands in the hard red winter wheat nursery were greatly reduced. The average yield for that test was 48 bushel/acre. The two offstation nurseries in Lake and Ravalli County had fair yields.

RESULTS:

The hard red winter Intrastate wheat nursery and the Western Regional Soft White Wheat nursery were severely injured by extreme temperatures that occured early February when wind chill factors exceeded 90 degrees below zero. Because of the severe damage to these two tests, the experiments were abandoned.

1989 Western Regional Hard Red Winter Wheat Nursery

Winter kill was 100% in three of the entries in this nursery. The variety UT 15751 and a survival rate of 87.5%, which was the highest for the nursery. Generally, the varieties had about 50% survival; however the test mean was 38.68%. Because of this stand reduction, there was a great reduction in yield as contrasted to previous years. The highest yielding entry was Idaho 381 at 82.5 bushels/acre. The average yield for the nursery was 48.63 bu/A. Test weights were reduced and the harvest date averaged about five days later than normal. Table 1.

1989 Lake County Offstation Winter Wheat Nursery.

Yields were very good for this nursery considering it was located on a recrop site. Stand losses due to winter kill were not consistent for varieties between replications. Location with in the nursery seemed to have as much effect on injury as did the variety. Yields range from 42.9 to 94.5 bu/A. Test weights were good with the mean being about 59.44 lbs/bu. Table 2.

1989 Ravalli County Offstation Winter Wheat Nursery

The yields from the Ross McIntyre farm in Ravalli Co. were some of the best that we recorded in that area. Rocky was the highest yielding entry with 60.9 bu/A. The lowest yielding entry was MT 88065. Test weights were fair with the average being 58.15 lbs/bu. Table 3.

FUTURE PLANS:

Winter wheat variety evaluations will continue with emphasis on selecting genetic material for dwarf smut resistance.

Table 1. Agronomic data from the 1989 Western Regional Hard Red Winter Wheat Nursery grown on the Northwestern Agricultural Research Center, Kalispell, MT.

Date planted: Sept. 16, 1988 Date harvested: Sept. 11, 1989

	VARIETY	Yield Bu/A	Test Wt Lbs/Bu			g % /1 Surviv		
ID 0381	ABERDEEN SEL	82.58	55.40	36.22	168.00	45.00	12.50	1.00
	WINTER WHEAT HYBRID	79.06	58.40	36.42		67.50	.00	.00
UT160719		77.54	55.15	34.25		75.00	.00	.00
	ABERDEEN SEL	76.60	56.80	35.83		72.50	.00	.00
	MNG/BEZ1	70.91	57.50	32.28		87.50	.00	.00
	HNL/USSR 2109-36	68.58	56.60	34.65		32.50	.00	.00
	LIND SELFRG/SRG (FIE	67.86	51.50	33.46		38.75	2.50	1.25
	LCD/FRD//NE 69559/WN	66.24	50.05	31.69		60.00	.00	.00
	CNN//7*LEE/TF/5/SM4/	65.81	56.20	30.51		45.00	.00	.00
	ABERDEEN SEL	63.25	55.25	26.97		55.00	.00	.00
	MNG/6/MT C61-9/BGR/5	62.89	53.50	29.13		82.50	.00	.00
	WINTER WHEAT HYBRID	62.15	55.15	29.53		48.75	.00	.00
	FRD/WNK//MT 6928/TR	60.90	53.75	36.22		72.50	.00	.00
	286011/ANDREWS	58.85	53.90	36.22		52.50	.00	.00
	ABERDEEN SEL	58.63	56.25	31.30		67.50	.00	.00
	ABERDEEN SEL	58.19	56.10	33.86		67.50	.00	.00
	PI173467/GNS/3/BNK//	57.78	50.30	28.35		32.50	.00	.00
CI 13844		57.56	55.45	37.99		60.00	8.75	1.50
	KHARKOF	56.85	54.55	42.32		72.50	91.25	5.25
	UT755079/CST56//TX65	56.40	52.30	32.48		42.50	.00	.00
	II60156/CI 14107//IT	55.49	51.60	35.63		37.75	18.75	4.25
	N7701501//V72044/CER	53.69	50.70	35.63		37.75	.00	.00
	HTN S/3/BEZ/CI 13438	52.56	51.90	31.30 1		17.75	.00	.00
	N7701501//V72044/CER	51.89	50.05	37.20 1		47.75	.00	.00
	HARD WHITE 1987 ML	49.25	52.85	28.35 1		23.75	.00	.00
NA007646		48.31	52.50	32.48 1		28.75	.00	.00
	N7000063/K71056//UT9	48.26	47.50	37.80 1		37.75	20.00	1.50
	CORVALLIS SEL	33.35	53.75	26.38 1		7.75	.00	.00
	BEZ1/MNG/3/HNL//IT/P	29.84	55.10	33.86 1	66.00	9.00	.00	.00
DR 8522	VORO/MNIM,85B-839	28.16	55.35	33.46 1		5.25	.00	.00
	HGL/RGR//A65249W-12-	26.05	55.75	28.15 1	66.50	12.75	.00	.00
	CORVALLIS SEL	25.42	53.05	28.35 1		17.50	.00	.00
DRCR8601	PMF//CNO S/GLL	24.95	52.50	31.30 1	64.25	5.75	.00	.00
	CORVALLIS SEL	21.30	54.25	24.61 1		12.75	.00	.00
	CORVALLIS SEL	18.68	50.30	24.61 1	64.75	12.75	.00	.00
DR830282	CORVALLIS SEL	17.47	53.60	22.44 1	65.25	16.25	.00	.00
DR832306	CORVALLIS SEL	2.16	.00	24.41 1	65.50	.00	.00	.00
DR830027	CORVALLIS SEL	1.00	.00	33.86	41.25	.00	3.75	1.00

^{*} Statistics and footnotes on next page

	Yield	Test Wt	Height Heading %			Lodging-		
VARIETY	Bu/A	Lbs/Bu	Inches	Date	Surviv	%	angle	
EXPERIMENTAL MEANS	48.63	49.61	32.01	160.23	38.68	4.04	.40	
F TEST FOR VAR. 2/		14.92	7.47			12.19		
C.V. 2: (S DF MEAN/MEAN) \$100 LSD (0.05)	12.24	1.12	5.06 4.63			107.29		

^{1/ %} Surviv = percentage of plot surviving winter kill

^{2/} F value for variety comparison, all F values were highly significant

Table 2. Agronomic data from the 1989 Offstation Winter Wheat Nursery grown on the Ed Wehrheim farm, Lake County, Moiese, MT.

Planted: September 13, 1988 Harvested: August 4, 1989

CI or		YIELD 1	EST WT	HEIGHT	%	LOI	GING
State #	VARIETY		LB/BU			ANGLE	%
					ET HIV		
CI 17846	MANNING	94.47	58.87	39.63	81.67	3.33	63.00
CI 17419	DAŴS	86.62	59.53	32.28	88.33	.00	.00
1T 88065	CST/VT1230//ID745101	77.19	59.23	40.55	50.00	.67	8.33
CI 17879		74.49	59.50	43.04	83.33	1.67	33.00
I 17909	LEWJAIN	73.87	58.57	39.63	61.67	.00	.00
1T 8003	TIBER	73.55	60.40	40.42	81.67	.00	.00
D 297	BLIZZARD	70.65	60.50	43.18	71.67	2.33	33.00
I 17735	NORSTAR	69.49	60.47	48.82	68.33	3.00	96.00
I 17860 I	NEELEY	68.99	59.10	36.61	45.00	2.00	64.67
I 17902	WINRIDGE	67.33	58.93	37.53	80.00	3.67	44.67
1 13670	WINALTA	66.93	59.00	46.33	55.00	4.33	83.33
1491533	NORWIN	65. 87	60.93	27.95	63.33	.00	.00
IT 88062	CST/VT1230//ID745101	65.67	59.20	41.60	81.67	1.67	35.00
I 17441	VDNA	63.82	60.73	35.17	43.33	.00	.00
I 17727	WESTON	61.88	62.20	41.86	65.00	1.33	56.67
T 8039		59.93	59.23	37.53	55.00	.00	.00
1491532		58.65	60.20	43.18	66.67	3.67	82.67
I 17594		57.43	56.87	37.14	40.00	.00	.00
I 8885 I	CHEYENNE	57.12	60.27	43.18	71.67	3.33	56.67
T 88064 I	CST/VT1230//ID745101	51.77	56.40	40.55	38.33	1.67	55.00
T 79125 I	UT755079/CST56//TX65	51.41	57.27	34.38	55.00	.00	.00
I 17844	REDWIN	51.05	59.90	42.13	40.00	.00	.00
I 13190 I	WARRIOR	49.70	59.47	43.44	60.00	2.67	58.00
I 15075	CENTURK	42.86	59.80	40.94	61.67	1.67	46.67
1.63	26.73		DIBATO	V10121		TECHS IN	
EXPE	RIMENTAL MEANS	65.03	59.44	39.88	62.85	1.54	34.03
F TES	ST FOR VAR.	1.59	4.01*	* 3.37**	1.32	2.53*	* 2.77*
C.V.	2:	14.55	1.09	6.19	21.11	60.00	56.83
LSD	(0.05)	26.93	1.84	7.03	37.76	2.63	55.05

** Indicates statistical significance at the .01 level of probability.

Special notes: When evaluating this agronomic data keep in mind that the nursery was recrop winter wheat, it was irrigated, and that winter kill was extensive in some varieties.

Table 3. Agronomic data from the 1989 Offstation Winter Wheat Nursery grown on the Ross McIntyre farm, Ravalli County, Stephens-ville, MT.

Planted: September 14, 1988	B Harvest	Harvested: August 1, 1989				
BOWN LOCK TOXAL SURVE AND		YTELSAV				
HARTETY	YIELD					
VARIETY	BU/A	LB/BU	INCHES			
01 47070 PROVV	/A 00	40.07	9 AT 1 13			
CI 17879 ROCKY	60.88					
ID 297 BLIZZARD	59.28					
CI 17846 MANNING	58.30					
MT 8003 TIBER						
CI 17735 NORSTAR	50.70					
CI 17441 VONA	49.72					
CI 17909 LEWJAIN	49.30					
CI 17419 DAWS	48.72					
MT 79125 UT755079/CST56//TX65 CI 17727 WESTON	47.93 47.25					
		59.33				
	47.00					
CI 13190 WARRIOR PI491532 CREE	46.93					
CI 13670 WINALTA	46.55					
	45.98					
PI491533 NORWIN						
MT 88062 CST/VT1230//ID745101	44.05	55.77	28.08			
CI 17844 REDWIN						
MT 8039 JUDITH						
CI 17860 NEELEY	42.88					
	42.43					
CI 17594 STEPHENS						
	40.67					
MT 88065 CST/VT1230//ID745101	39.15					
	28/535	125779413	7757			
EXPERIMENTAL MEANS	47.57	58.15	28 44			
F TEST FOR VAR.	.73					
C.V. 2: (S DF MEAN/MEAN) \$100						
LSD (0.05)		1.81				
Lan (0.03)	10.00	1.01	0.07			

^{**} Indicates statistical significance at the .01 level pf probability.