TITLE:

Winter Wheat

PROJECT:

Small Grains Investigations MS 756

YEAR:

1968

PERSONNEL:

Leader:

Vern R. Stewart

Cooperators:

J. R. Welsh, Charles McQuire

LOCATION:

Northwestern Montana Branch Station and several off station locations throughout Montana which will be identified in the

manuscript

DURATION:

Indefinite

OBJECTIVES:

1. To obtain the information necessary for making varietal recommendations and evaluating new varieties and selections.

2. To conduct a breeding program in northwest Montana designed to produce high yielding varieties with particular emphasis on the acceptable quality and resistance for dwarf bunt and stripe rust. Other agronomic characteristics such as straw strength, winter hardiness etc., will be evaluated in this program.

SIGNIFICANT FINDINGS:

Crest is 129% of Westmont over a four year period, Delmar does exceed Crest in yield for the same period.

PI 178383 + Westmont crosses provide a high degree of resistance to

the dwarf smut.

Wa 4995 was the high yielding entry in the white wheat nursery.

In all locations summarized, Omar was high in yield of the white wheats and McCall the highest for hard red varieties.

FUTURE PLANS:

Plans for 1968-69 include regular yield nurseries and assistance in the overall state breeding program.

MATERIALS AND METHODS:

Standard nursery procedures were used in all of the variety testing programs. A randomized block design was used having four to six replications. Data obtained were: yield; plant height; test weight; disease and lodging. Nurseries grown were: Intrastate Winter Wheat Nursery at the Northwestern Montana Branch Station in Field E-4; Western Regional Hard Red Winter Wheat Nursery grown on the L. B. Claridge farm, northwest of Kalispell in a dwarf bunt area; Uniform White Wheat Nursery grown at the Northwestern Montana Branch Station in Field E-4. The off station nurseries were located in Ravalli, Missoula, Lake, Sanders and Mineral Counties. A nursery on the Lance Claridge farm northwest of Kalispell was grown to increase seed supply of breeding lines.

MATERIALS AND METHODS (con't):

Herbicide applications were made for weed control in the off station locations in early spring. Herbicide applications at Claridges and on the station were made in the fall. Where necessary follow-up applications were made in the spring to control any spring germinating weeds. The herbicide used for weed control was bromoxynil at a rate of 3/8 of a pound per acre. Plots were harvested with a power harvester.

RESULTS AND DISCUSSIONS:

Intrastate Nursery

Winter wheat was about average in yield in 1968 at the Northwestern Montana Branch Station. The highest yielding entry in the nursery was McCall at 79.0 bu/acre. Several varieties were found to be significantly lower in yield than Delmar the check variety. Dwarf Smut infection was not a factor in this nursery in 1968. Stripe rust readings were made and recorded. Lodging was severe in all varieties, which made harvest difficult. Test weights were above average. Table 1 shows tabulation of the data in this study.

In Table 2 is shown a summary of the data for 10 years at the Northwestern Montana Branch Station. Only two varieties in this years study have been grown in this study for 10 years. Over the 10 year period Cheyenne has out yielded the variety Westmont. This is due primarily to the dwarf bunt incidence that occurs in this region at times. Over a four year period Crest is 129% of Westmont. Over a four year period this variety has yielded 54.8 bu/acre as contrasted to Westmont with 42.5 bu/acre and Cheyenne 52.9 bu/acre. Delmar is the only variety that has a higher yield than Crest over a four year period.

Western Regional Hard Red Winter Wheat Nursery

Yield data was not significant in this nursery in 1968. Dwarf smut was light in this study. Some of the Westmont/PI 178383 back crosses show complete immunity to dwarf bunt. Crest has a O reading in 1968. Cheyenne a susceptible variety has a reading of .75%. Complete tabulation of these data are found in Table 3.

Test weights are excellent in this nursery.

Uniform White Wheat Nursery

High yields were obtained from the white wheat nursery grown on the station. Lodging resistance was very good in most of the white entries except Moro which tends to have weak straw. Golden and Karkoff in this study had high incidence of lodging. The high yielding entry was Wa 4995 with 98.83 bu/acre and was significantly higher in yield than Nugaines, the check. Test weights were below standard in many of the entries. (Table 4)

Table __4 . Agronomic data from the uniform white winter wheat nursery, North-western Montana Branch Station, Field No. E-4, 1968. Experimental design - random block, 4 replications.

Seeding Date: September 17, 1967 Harvest Date: August 13, 1968 Size of Plot: 16 sq. ft.

		Yield	Heading	Plant	Test Wt	Stripe	Rust	Lodg	ing _
Variety	Number	bu/a	Date	Height	1bs/bu	Sev.	Type	%	Sev.
N10/B11/P14/14/53	-		/ / /	00 00	r/ 0	0 05	2.00	77 50	2.00
101	4995	98.83*		39.00	56.8	0.25	2.00	17.50	4.00
Suwon 92 x 40mar	4966	98.08	6/20	43.25	60.0	0.25	2.00	25.00	4.00
Nord Desprey/2	/0770	00 05	6.00	36.50	54.7	0.25	2.00	12.50	1.25
Sel. 101	63112	90.05	6/18	5y-9,11-1-2000		3.25	6.25	45.00	5.25
Brevor	12385	87.90	6/20	44.75	61.9	2027	0.2)	47.00	102)
(14/53/0din)CI	I ARR	86.68	6:/20	40.50	56.2	0.00	2.00	10.00	1.25
13431	4877			45.00	60.1	7.50	7.25	28.75	2.75
Burt	12696	86.50		47.75	57.3	1.50	3.50	66.25	5.75
Moro 1/	13740	86.33		34.00	59.5	1.75	3.25	10.00	1.50
Nugaines1/	13968	85.83			59.5	1.00	3.00	11.25	1.25
Suwon 92/4 Burt	4984	85.45		34.25	57.1	9.00	7.25	13.75	1.75
Gaines	13448	84.35		34.00	60.7	20.00	7.50	58.75	4.75
Elgin	11755	80.45	6/21	50.00	61.0	4.00	3.25	27.50	3.50
PI 178383/40mar	88	80.18	6/21	51.00	District of Alberta		7.75	48.75	5.75
Omar	13072	73.85	6/21	51.75	59.0	27.50	7.75	66.25	5.25
Triplet	5408	71.32	6/16	51.50	62.4	12.50	8.00	67.50	7.50
Golden	10063	70.35	6/20	54.00	57.8	20.00	4.75	73.75	7.75
Kharkof	1442	58.52	6/19	49.25	61.3	6.75	4.12	13.13	1.1)

1/ check variety
 variety yielding significantly more than the check (.05)

x	82.8
S.E.X	4.6
L.S.D.(.05)	12.96
C.V.%	5.50

Analysis of Variance

Source	D.F.	H.S.	F.
Replications	3	163.3	1.97
Varieties	15	426.3	5.14*
Error	45	82.9	
Total	63		