



**On-Farm Cereal Variety and Advanced Breeding Line  
Testing across Montana for Environment Specific  
Cultivar Recommendations:**



**Winter Wheat Off-Station Variety Performance, Loma, MT**

**Principal Investigator:**

Peggy Lamb, Senior Research Scientist, Northern Ag Research Center, Havre

**Project Personnel:**

Sue Mondal, Breeder/Geneticist, Winter Wheat, Bozeman  
Andrew Lehnerz, Research Associate, Winter Wheat, Bozeman  
Arpit Gaur, Post-Doc, Winter Wheat, Bozeman  
Eleri Haney, Research Associate, Havre  
Reese Whitehead, Research Assistant II, Havre  
Tyler Lane, Chouteau County Extension

**Cooperators:**

Terry McKeever, Lyle McKeever and Kendrick McKeever, Landowners, Loma

**Objectives:**

Commercially available winter wheat varieties and advanced breeding lines were evaluated for agronomic performance and fit at on-farm locations across the state of Montana. Sites chosen for the research considered the environment, growing conditions and soil types, and represent the major land areas for producers in those regions served by Northern Agricultural Research Center. The Chouteau County location, between Big Sandy and Loma, has been used for various trials since 1998.

**Methods:**

The uniform off-station winter wheat variety performance trial was seeded into chemical fallow ground during 2025. The trial consisted of 25 entries seeded in replicated, three-row, 22-foot plots on a 12-inch row spacing, utilizing a self-propelled cone seeder with Atom Jet paired row openers. All plots were trimmed to a harvest length of approximately 17 feet with a three-point rototiller. Plant height was measured from the soil surface to the top of the head, excluding awns, and percent sawfly cutting was visually estimated for each plot immediately prior to harvest. A 'Wintersteiger' small plot combine, funded in part by the Montana Wheat and Barley Committee, was used to harvest each three-row plot. Prior to measuring plot weight for yield determination, seed was either cleaned or weighed in-dirt as per protocols. Protein, test weight, and moisture content were determined on a clean sample using a Foss Infratec 1241 near infrared analyzer. Falling number was determined using a Perten FN1700 according to the FGIS Directive 9180.38. Other variables specific to each individual trial are listed with the current year's data tables.

Please note that research trial seed yield results recorded under wheat stem sawfly pressure are likely much higher than a producer should expect. Small plot variety trials are managed to assess maximum yield potential and are harvested in such a way that all stems and heads are picked up by the combine, regardless of lodging or cutting due to wheat stem sawfly. Pickup guards coupled with an exceptionally slow ground speed and a low cutting height help researchers collect all heads in order to assess seed

yield potential. If you are a producer in a wheat stem sawfly environment, although hollow stemmed varieties may be high yielding in research trials in your area, we strongly recommend against growing those hollow stemmed varieties. Please be aware that if you seed hollow stemmed varieties with wheat stem sawfly present, you are only creating a breeding ground for future generations of sawfly in your area and not helping combat the pest population.

Results:

This report contains both single-year and long-term data summaries limited to the most recent ten years. It should be noted that the 2025 data table in this report represents varietal performance for a single crop year at a single location, therefore cannot be considered representative of performance expected when differing conditions due to location, year and management are imposed. By itself, 2025 data shall not constitute in any form a recommendation for or against any variety or breeding line included.

Numerous heavy rain events occurred following maturity, negatively influencing seed quality. In conjunction with wheat stem sawfly infestation, this site also had hessian fly pressure. At the Loma winter wheat site, seed yields averaged over 68 bu/ac (Table 1). Montana State University breeding line 'MTCS22127' was the highest yielding entry at just under 78 bu/ac. 'Bobcat', 'Brawl', 'Fourosix', 'MT Meadowlark', 'Northern', 'SY Monument', and four MSU breeding lines produced yields statistically equal to that of the highest yielding entry. Test weights of all entries averaged just under 57 lb/bu. Wheat stem sawfly cutting was severe in the winter wheat trials averaging just under 50 percent cut and lodged. Yield, test weight, protein, falling number, plant height, and sawfly cutting data for the 2025 Loma dryland winter wheat trial are summarized in Table 1.

Comparable averages are calculated using a standard check variety when not all entries are present in a specific trial for all years. Variety means are adjusted by multiplying the actual check mean by the ratio of the individual variety mean compared to the check mean for the same years as tested. All varieties are then directly comparable to each other when in the same nursery. A minimum of three years of data is necessary to be included in the comparable average calculation. Ten-year comparable averages for seed yield and test weight at Loma are summarized in Table 2, while ten-year comparable averages for protein content and sawfly cutting are summarized in Table 3. Based on the comparable average calculations, Bobcat is the highest yielding variety at the Loma site and has the least amount of wheat stem sawfly cutting across the years.

Summary:

Due to late seeding, stands were not fully established prior to winter. Precipitation was above average during late May and June, resulting in good yield potential. However, continued rains after harvest maturity resulted in poor quality seed and low falling numbers.

This work has been strongly supported by producers in the Loma-Big Sandy area, and by the Northern Agricultural Research Center Advisory Board. With budget and other resources allowing, it is planned to continue off-station winter wheat variety and breeding line testing in this area. The Chouteau County location between Loma and Big Sandy has been hosting various trials for 27 years, beginning in 1998.

Recognition:

This research would not have been possible without the assistance of the following seasonal employees: David Bischoff, Baylor Davis, Ty Golie, Simone Gomes, Brady Kueffler, Lirio McKenzie, Teresa Miller, Jason Rice, and Lily Smith.

**TABLE 1. Dryland Fallow Winter Wheat Cultivar Evaluation Nursery Grown Off-Station at McKeever Farm & Seed, Inc., Loma. Northern Agricultural Research Center. Havre, Montana. 2025. (Exp# 25-3853-WW)**

ID	1/ YIELD bu/ac	2/ TEST WT lb/bu	3/ PROTEIN %	FN seconds	4/ PLNT HT inches	SAWFLY %
AAC Coldfront	60.0	56.2	<b>14.6</b>	115	28.7	81.7
AAC Wildfire	63.5	55.9	14.2	173	<b>30.3</b>	70.0
Bobcat	<b>73.7</b>	57.7	14.3	286	26.4	<b>10.0</b>
Brawl CL Plus	<b>75.9</b>	57.8	14.4	219	<b>30.0</b>	31.7
CS Bridger CLP	66.9	56.2	13.8	80	27.6	41.7
Flathead	67.1	57.1	13.8	107	29.2	60.0
Fortress	67.1	55.7	13.5	194	27.3	43.3
FourSixx	<b>77.5</b>	56.3	13.9	265	29.0	65.0
Keldin	63.0	57.0	<b>14.5</b>	169	28.1	53.3
Loma	60.8	56.0	<b>14.5</b>	88	26.5	65.0
MT Meadowlark	<b>68.9</b>	57.1	13.9	280	28.9	30.0
MT WarCat	65.9	57.2	14.0	<b>384</b>	28.4	23.3
Northern	<b>73.6</b>	55.6	13.9	90	<b>30.5</b>	53.3
StandClear CLP	63.0	56.9	14.3	163	29.5	46.7
SY Monument	<b>69.3</b>	55.5	13.2	284	29.0	56.7
Warhorse	54.1	57.1	<u>15.0</u>	<b>338</b>	28.5	21.7
Yellowstone	60.4	55.8	<b>14.7</b>	117	<b>31.1</b>	56.7
MT2142	<b>68.6</b>	56.5	14.0	245	27.0	75.0
MT2270	63.4	57.7	13.6	103	27.8	53.3
MTAX21203	<b>77.2</b>	57.1	13.1	207	29.5	68.3
MTAX22120	67.3	56.4	13.8	290	28.3	31.7
MTCS22127	<b>77.9</b>	56.4	13.9	270	<b>30.0</b>	53.3
MTCS2264	<b>70.1</b>	<b>58.7</b>	<b>14.8</b>	179	27.0	66.7
MTS2286	<b>76.8</b>	56.6	14.1	212	28.7	35.0
MTV2164	<b>77.3</b>	56.0	13.6	294	<b>31.8</b>	51.7
EXPERIMENTAL MEANS	68.4	56.7	14.1	206.1	28.8	49.8
LSD (0.05)	9.6	0.5	0.5	48.6	1.8	24.1
C.V.%	8.5	0.5	2.0	14.4	3.8	29.5
P-VALUE (Varieties)	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001

1/ Volumetric yields are based on plot weights adjusted to uniform 13 percent grain moisture and 60 lbs/bu as the standard test weight for wheat.

2/ Protein values are adjusted to 13 percent grain moisture.

3/ FN is the falling number value reported in seconds adjusted to 14 percent flour moisture.

4/ Sawfly rating is reported as the percentage of cut stems.

**Bold** indicates the highest or lowest value within a column (whichever is most desirable for the specific characteristic).

**Bold** indicates values equal to the underlined value within a column based on Fisher's protected LSD (P=0.05).

Management Information (25-3853-WW)

Seeding Date:	October 8, 2024
Harvest Date:	August 19, 2025
Fertility:	100-20-10-10 side banded
System:	No Till
Herbicide:	Brox M 24 oz; Opensky 16 oz
Insecticide:	none
Previous Crop:	Chemical Fallow-Spring Wheat
Precipitation:	8.74" April 1 to harvest maturity

TABLE 2. Ten-Year Yield and Test Weight Summary on Selected Entries from Dryland Fallow Spring Wheat Variety Nurseries Grown Off-Station at McKeever Farm & Seed, Inc., Loma. Northern Agricultural Research Center. Havre, Montana. 2016-2025. (Exp# 3853-WW)

2/ VARIETY	3/ No. of YEARS TESTED	1/ YIELD (Bushels Per Acre)						5/ 10-YR COMP. AVE. YIELD	TEST WEIGHT (Pounds Per Bushel)								
		3/ AVE. for YEARS TESTED					4/ % of CHECK		3/ AVE. for YEARS TESTED					5/ 10-YR COMP. AVE. TEST WT			
		2021	2022	2023	2024	2025	TEST WT	2021	2022	2023	2024	2025	TEST WT				
BOBCAT (++) (sawfly tol)	9	31.8	41.5	74.8	64.5	73.7	59.5	109.7	58.4	54.3	56.7	60.7	55.6	57.7	58.8	103.0	58.5
BRAWL CLP (++) (CL)	9	36.3	36.1	75.8	55.7	75.9	57.0	105.1	56.0	55.2	59.1	62.7	54.2	57.8	59.5	104.3	59.3
NORTHERN (+)	10	28.1	34.7	88.3	49.1	73.6	55.1	103.5	55.1	52.8	55.1	60.5	52.9	55.6	56.9	100.2	56.9
SY MONUMENT (P+)	9	26.7	34.0	76.1	57.1	69.3	55.6	102.4	54.5	52.2	58.7	59.4	51.9	55.5	57.5	100.7	57.2
FOUROSIX (++)	9	29.9	32.4	74.9	53.0	77.5	54.3	100.0	53.3	51.9	57.6	58.8	52.3	56.3	57.5	100.7	57.2
YELLOWSTONE (+)	10	27.5	40.2	84.6	61.5	60.4	53.2	100.0	53.2	52.5	56.0	58.5	53.2	55.8	56.8	100.0	56.8
KELDIN (P+)	10	27.2	35.9	77.4	53.5	63.0	53.2	99.9	53.2	53.5	55.3	59.8	52.5	57.0	57.5	101.2	57.5
BRIDGER CL PLUS	4		38.4	82.9	57.8	66.9	61.5	99.8	53.1		57.7	60.0	53.3	56.2	56.8	101.7	57.8
FLATHEAD (++)	8	29.3	31.4	78.1	59.2	67.1	54.4	97.7	52.0	54.4	58.7	60.5	53.8	57.1	58.3	102.8	58.4
LOMA (++)	10	30.5	36.5	85.8	54.4	60.8	51.7	97.1	51.7	53.0	56.4	58.5	53.6	56.0	57.0	100.4	57.0
STANDCLEAR CLP	8	29.1	40.1	80.8	58.3	63.0	54.1	97.0	51.6	54.5	58.0	60.5	55.0	56.9	58.5	103.1	58.6
MT WARCAT (++)	6	26.9	39.2	88.0	43.8	65.9	53.5	96.2	51.2	52.7	56.4	58.7	54.2	57.2	56.5	101.0	57.4
AAC WILDFIRE (++)	7	22.5	35.6	78.2	49.1	63.5	52.2	94.4	50.2	51.4	56.1	58.4	52.4	55.9	56.1	99.4	56.5
WARHORSE (+) (sawfly tol)	10	28.1	27.7	73.6	50.0	54.1	45.1	84.7	45.1	52.2	56.8	58.7	54.0	57.1	57.6	101.4	57.6
MEANS (For Entries Listed)		28.8	36.0	80.0	54.8	66.8			52.8	53.1	57.0	59.7	53.5	56.6			57.6

6/ Growing Season Precipitation (in.)	n/a	1.60	8.43	n/a	8.07	6.53
Soil PAW (in.) to SD @ Planting	7.94	7.12	n/a	7.34	7.92	7.75
Total Plant Available Water (in.)	n/a	8.72	n/a	n/a	15.99	13.81
Soil NO <sub>3</sub> (lbs.) to SD at Planting	97	299	n/a	114	33	137
SD (Sampling Depth in Inches)	45	48	n/a	45	44	47
Fertilizer Applied	(# N)	125	125	89	92	100
	(# P <sub>2</sub> O <sub>5</sub> )	20	20	14	50	20
	(# K <sub>2</sub> O)	10	10	7	7	10
	(# S)	10	10	7	7	10

Check Variety is Yellowstone.

1/ See MCES Bulletin 1098 or the Plant Sciences & Plant Pathology website at <http://plantsciences.montana.edu/> for evaluation of other important variety performance characteristics to include protein, quality, winter hardiness, disease resistance, etc. before making cultivar selection decisions.

2/ P = Private Variety, + = Protected Variety, ++ = PVP Title 5 Pending, CL = Clearfield Line, HW = Hard White.

3/ Only the most recent 5 years shown, but summary calculations include all years noted.

4/ Percent of Yellowstone yield or test weight for the same data years as those in which a given entry was tested.

5/ 10-Yr Comparable Average =  $(x/y) * z$  where x = average yield or test weight of a given entry for years tested, y = average yield or test weight for Yellowstone for the same years, and z = 10-Yr average yield or test weight for the check variety Yellowstone.

6/ April 1 to 14 days prior to harvest

TABLE 3. Ten-Year Protein and Sawfly Summary on Selected Entries from Dryland Fallow Spring Wheat Variety Nurseries Grown Off-Station at McKeever Farm & Seed, Inc., Loma. Northern Agricultural Research Center, Havre, Montana. 2016-2025. (Exp# 3853-WW)

2/ VARIETY	3/ No. of YEARS TESTED	1/ PROTEIN % (Adjusted to 13% grain moisture)										SAWFLY RATING (% of cut and lodged stems)									
		3/ AVE. for YEARS TESTED					4/ % of CHECK PROTEIN		5/ 10-YR COMP. AVE. PROTEIN		3/ AVE. for YEARS TESTED					4/ % of CHECK PROTEIN		5/ 10-YR COMP. AVE. SAWFLY			
		2021	2022	2023	2024	2025	TESTED	PROTEIN	CHECK	PROTEIN	2021	2022	2023	2024	2025	TESTED	SAWFLY	CHECK	PROTEIN		
BOBCAT (++)(sawfly tol)	10	14.8	15.0	12.3	14.5	14.3	13.9	99.4	14.2	14.2	15.9	1.0	2.3	18.3	10.0	10.0	16.6	16.6	9.0		
WARHORSE (+)(sawfly tol)	8	15.1	15.1	12.4	14.7	15.0	14.8	103.6	14.8	14.8	16.9	7.0	2.3	51.7	21.7	15.1	27.8	27.8	15.1		
MT WARCAT (++)	9	15.3	14.9	12.5	15.3	14.0	14.4	103.8	14.8	14.8	52.5	8.3	31.7	90.0	23.3	35.3	57.3	57.3	31.0		
LOMA (++)	10	15.1	15.1	11.7	14.2	14.5	14.4	101.1	14.4	14.4	68.0	13.3	35.0	96.0	65.0	38.7	71.5	71.5	38.7		
STANDCLEAR CLP	8	14.5	14.5	12.3	13.7	14.3	13.7	99.3	14.2	14.2	57.5	16.7	58.3	85.0	46.7	46.1	72.6	72.6	39.3		
BRAWL CLP (++)(CL)	10	14.7	14.6	12.1	14.5	14.4	14.0	100.4	14.3	14.3	61.1	45.0	55.0	66.7	31.7	45.1	75.4	75.4	40.8		
FLATHEAD (++)	6	15.0	14.7	11.6	13.8	13.8	13.7	99.0	14.1	14.1	62.6	11.7	73.3	93.3	60.0	53.5	84.2	84.2	45.6		
AAC WILDFIRE (++)	7	15.2	14.6	11.8	14.1	14.2	13.8	100.6	14.4	14.4	79.4	58.3	36.7	90.0	70.0	52.2	86.3	86.3	46.7		
SY MONUMENT (P+)	7	13.6	14.1	11.3	13.1	13.2	12.9	92.3	13.2	13.2	74.8	51.7	40.0	86.7	56.7	52.3	87.4	87.4	47.3		
NORTHERN (+)	8	15.5	15.5	11.1	14.5	13.9	14.6	102.0	14.6	14.6	57.9	65.0	70.0	90.0	53.3	50.1	92.6	92.6	50.1		
BRIDGER CL PLUS	4		14.8	11.4	14.5	13.8	13.6	99.3	14.2	14.2	55.0	88.3	93.0	41.7	69.5	98.2	98.2	53.2			
YELLOWSTONE (+)	10	14.6	14.8	11.5	14.0	14.7	14.3	100.0	14.3	14.3	81.4	56.7	73.3	96.3	56.7	54.1	100.0	100.0	54.1		
KELDIN (P+)	8	14.3	15.4	12.0	14.5	14.5	14.2	99.6	14.2	14.2	76.8	73.3	41.7	78.3	53.3	55.2	102.0	102.0	55.2		
FOUROSIX (++)	5	14.9	14.8	11.8	15.0	13.9	13.9	99.7	14.2	14.2	84.1	53.3	66.7	85.0	65.0	63.3	105.7	105.7	57.2		
MEANS (For Entries Listed)		14.8	14.9	11.9	14.3	14.2			14.3	14.3	60.7	36.9	48.2	80.0	46.8				41.7		

6/ Growing Season Precipitation (in.)	n/a	1.60	8.43	n/a	8.07	6.53
Soil PAW (in.) to SD @ Planting	7.94	7.12	n/a	7.34	7.92	7.75
Total Plant Available Water (in.)	n/a	8.72	n/a	n/a	15.99	13.81
Soil NO <sub>3</sub> (lbs.) to SD at Planting	97	299	n/a	114	33	137
SD (Sampling Depth in Inches)	45	48	n/a	45	44	47
Fertilizer Applied	(# N)	125	125	89	92	100
	(# P <sub>2</sub> O <sub>5</sub> )	20	20	14	50	20
	(# K <sub>2</sub> O)	10	10	7	7	10
	(# S)	10	10	7	7	10

Check Variety is Yellowstone.

1/ See MCES Bulletin 1098 or the Plant Sciences & Plant Pathology website at <http://plantsciences.montana.edu/> for evaluation of other important variety performance characteristics to include protein, quality, winter hardiness, disease resistance, etc. before making cultivar selection decisions.

2/ P = Private Variety, + = Protected Variety, ++ = PVP Title 5 Pending, CL = Clearfield Line, HW = Hard White.

3/ Only the most recent 5 years shown, but summary calculations include all years noted.

4/ Percent of Yellowstone protein or sawfly for the same data years as those in which a given entry was tested.

5/ 10-Yr Comparable Average =  $(x/y) * z$  where x = average protein or sawfly rating of a given entry for years tested, y = average protein or sawfly rating for Yellowstone for the same years, and z = 10-Yr average protein or sawfly rating for the check variety Yellowstone.

6/ April 1 to 14 days prior to harvest