

Project Title: Evaluation of Wild Oat Herbicides in Spring Wheat

Project Leaders: Bob Stougaard

Project Personnel: Qingwu Xue

Objective: To evaluate reduced rate herbicide performance in spring wheat

Results:

Seven herbicides were applied at their respective label 1X and half-label 1/2X rates to evaluate the consistency of wild oat control in spring wheat. The experiment design was a randomized complete block with 3 replications. Scholar spring wheat was planted on April 28, 2006 at a seeding rate of 75 lb/ac in 7" rows to a depth of 1.5 inches. Wild oat was planted within each plot at a density of 25 seeds per square foot. Herbicides were applied on May 24, 2006 using a backpack sprayer with Teejet XR11002 nozzles in 20 GPA. Spring wheat and wild oat were at 4- and 3-leaf stages, respectively, at the time of application.

Crop injury was initially observed with most herbicides, particularly at the labeled rate, but only Silverado and Everest resulted in noticeable injury as the season progressed. However, the effects were transitory and injury was minimal by 8 weeks after application.

Except for Puma at 1/2X rate, all herbicides provided excellent wild oat control (>93%) regardless of the rate. However, herbicide rate did impact wild oat biomass, which was generally greater when herbicides were applied at the half rate as compared to the labeled rate. Correspondingly, yields tended to be slightly less for the half-labeled rate treatments. Herbicide rate did not affect dockage, grain moisture or test weight.

Summary:

All herbicides treatments provided excellent wild oat control except Puma at the reduced rate. Application of herbicides at half-label rates resulted in slightly lower yields and more wild oat biomass than at label rates.

Table 1. Effects of reduced herbicide rates on crop injury and wild oat control at Kalispell, MT during 2006.

Treatment	Label rate (1X, lb ai/ac)	Crop injury (%)								Wild oat control (%)					
		5/30/06		6/5/06		7/6/06		8/3/06		6/5/06		7/5/06		8/3/06	
		1X	1/2X	1X	1/2X	1X	1/2X	1X	1/2X	1X	1/2X	1X	1/2X	1X	1/2X
Achieve	0.1800	1.7	0.0	0.0	0.0	0.0	1.7	0.0	0.0	91.7	83.3	100.0	98.3	99.3	99.0
Axial	0.0540	0.0	0.0	0.0	3.3	0.0	1.7	0.0	0.0	91.7	91.7	100.0	95.0	100.0	99.0
Everest	0.0262	6.7	6.7	11.7	3.3	3.3	1.7	1.7	0.0	83.3	76.7	91.7	88.3	100.0	100.0
Silverado	0.0028	8.3	6.7	16.7	13.3	5.0	1.7	0.0	0.0	81.7	83.3	99.0	83.3	99.3	95.0
Hoelon	1.0000	8.3	3.3	0.0	0.0	1.7	0.0	0.0	0.0	88.3	80.0	100.0	93.3	100.0	98.3
Puma	0.0830	8.3	1.7	0.0	0.0	0.0	0.0	0.0	0.0	91.7	83.3	96.7	76.7	96.3	85.0
Discover	0.0500	1.7	0.0	1.7	0.0	0.0	0.0	0.0	0.0	78.3	81.7	100.0	93.3	99.0	93.3
Control		0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
LSD (0.05)	Herbicide (A)	5.7		3.5		2.3		NS		NS		4.1		1.3	
	Rate (B)	NS		1.9		NS		NS		NS		2.2		0.7	
	AxB	NS		4.9		NS		NS		NS		5.8		1.9	

Data from the control plots were excluded from the analysis.

Table 2. Effects of reduced herbicide rates on wild oat (WO) biomass, dockage, and spring wheat yield, grain moisture and test weight at Kalispell MT, during 2006.

Treatment	Label rate (1X, lb ai/ac)	WO Biomass (g/m ²)		Dockage (%)		Yield (bu/ac)		Grain moisture (%)		Test weight (lb/bu)	
		7/7/06		8/8/06		8/8/06		8/8/06		8/8/06	
		1X	1/2X	1X	1/2X	1X	1/2X	1X	1/2X	1X	1/2X
Achieve	0.1800	4.7	0.0	0.6	0.8	62.7	57.0	11.6	10.9	63.3	62.6
Axial	0.0540	0.0	5.9	1.2	0.8	63.5	56.7	14.0	11.4	62.3	62.6
Everest	0.0262	5.5	27.7	0.7	0.7	54.1	56.7	11.0	11.3	62.1	62.9
Silverado	0.0028	7.2	38.7	0.7	0.9	55.3	54.6	11.5	12.0	62.3	62.4
Hoelon	1.0000	0.0	4.9	0.9	0.8	58.8	58.8	11.8	11.1	62.4	62.7
Puma	0.0830	6.7	21.6	0.9	1.3	61.3	51.3	12.1	10.3	62.7	62.7
Discover	0.0500	0.0	9.1	0.9	1.2	66.0	61.4	12.9	12.2	62.8	62.9
Control		376.1	434.7	7.9	8.9	35.3	37.4	12.0	12.5	62.0	63.0
LSD (0.05)	Herbicide (A)	10.2		NS		5.1		NS		NS	
	Rate (B)	5.5		NS		2.7		NS		NS	
	AxB	14.5		NS		NS		NS		NS	

Data from the control plots were excluded from the analysis.